

Generalizations of two-stack-sortable permutations

A Dissertation

Presented to

The Faculty of the Graduate School of Arts and Sciences

Brandeis University

Department of Mathematics

Professor Ira Gessel, Advisor

In Partial Fulfillment

of the Requirements for the Degree

Doctor of Philosophy

by

Dapeng Xu

September, 2002

Acknowledgment

My deepest gratitude goes to my advisor, Ira Gessel, without whom this work would not be possible; I would like to thank him for his generosity in sharing his insight and time with me, and his constant patience and encouragement. It was and will always be my pleasure to work with him.

I would like to thank Daniel Ruberman, Kiyoshi Igusa, Alan Mayer and Mark Adler for teaching me a lot of mathematics. I would like to thank Susan Parker for helping me to be a better teacher.

I'd also like to thank the Mathematics Department faculty and staff for making my life here more enjoyable.

ABSTRACT

Generalizations of two-stack-sortable permutations

A dissertation presented to the faculty of
the Graduate School of Arts and Sciences of
Brandeis University, Waltham, Massachusetts

by Dapeng Xu

In this thesis, we apply the stack sorting operator to r -permutations and construct the functional equation for the generating function of two-stack-sortable k -tuple r -permutations counted by descents by using a factorization similar to Zeilberger's. We solve the functional equation and give explicit formulas for the number of two-stack-sortable r -permutations.

Contents

Chapter 1. Introduction	3
1.1. The generalization	5
1.2. Stack-sortable r -permutations	8
Chapter 2. Two-stack-sortable r -permutations	15
2.1. The functional equations	15
2.2. The solution to the functional equations	20
2.3. The number of two-stack-sortable r -permutations with descents	24
2.4. The number of two-stack-sortable r -permutations	31
Chapter 3. Parallel results	35
3.1. Another approach	35
3.2. Connection to Zeilberger's functional equation	39
Chapter 4. Further results	41
4.1. A generalization of the stack-sorting operation on r -permutations	41
4.2. A characterization of t -stack-sortable permutations	48
4.3. A modification on the functional equations.	52
Bibliography	55

CHAPTER 1

Introduction

The operation of stack sorting was first studied by Knuth [15, p. 239]. He described the operation by a railroad cars swapping algorithm. Later West described this operation in terms of a simple card game [27].

The operation uses a stack and can be described as follows. Let π be a word with distinct letters in the alphabet $\{1, 2, 3, \dots\}$. We call π the input of the operation. At the i th step, we compare the i th letter a of the word π with the top letter on the stack (if any). If a is smaller or the stack is empty, we put a on the top of the stack. If a is bigger, we move the top letter on the stack to the output and then again compare a with the new top letter on the stack (if any). We repeat this until we put a onto the stack. When the input is empty, we move all the letters on the stack (if any) to the output in order from top to bottom.

West also gave a recursive definition of the stack sorting operation. [27, 26].

DEFINITION 1.0.1. Let $\pi = a_1 a_2 \cdots a_n$ be a word in the alphabet $\mathbb{P} = \{1, 2, \dots\}$, having all its letters distinct. If $n = 0$, let $S(\pi)$ be the empty word. Otherwise, let $S(\pi)$ be obtained by permuting the letters of π as follows: if $m = \max(a_1, a_2, \dots, a_n)$ and $\pi = \pi_l m \pi_r$, then

$$S(\pi) = S(\pi_l) S(\pi_r) m.$$

It is easy to see that this recursive definition of the stack sorting operation is equivalent to the description in terms of stacks. For a given word $\pi = a_1 a_2 \cdots a_n$ in

the alphabet $\mathbb{P} = \{1, 2, \dots\}$, having all its letters distinct, we can always write it as $\pi = \pi_l m \pi_r$ where $m = \max(a_1, a_2, \dots, a_n)$. According to the description, since m is bigger than any letter in π_l , all the letters in π_l will be moved to the output before we can put m onto the stack. Also since m is bigger than any letter in π_r , m will not be removed from the stack until all the letters in π_r go through the stack. That is exactly what the recursive definition says.

The problem of stack sorting was then generalized and researched in a number of ways. Among these, the problems of enumerating t -stack-sortable permutations interest most people. We say that a permutation π of $[n]$ is a t -stack-sortable permutation if $S^t(\pi)$ is the identity permutation, and that π is exactly t -stack-sortable if π is t -stack-sortable but not $(t-1)$ -stack-sortable. Knuth [15, p. 239] proved that the number of 1-stack-sortable permutations of $[n]$ is the Catalan number $C_n = \binom{2n}{n}/(n+1)$. West [26, 27] proved that all permutations of $[n]$ are $(n-1)$ -stack-sortable, that $(n-2)!$ are exactly $(n-1)$ -stack-sortable, and that $\frac{7}{2}(n-2)! + (n-3)!$ are exactly $(n-2)$ -stack-sortable.

Characterizations of 1-stack-sortable permutations and 2-stack-sortable permutations were also given using the following notion of pattern avoidance. Let $\pi_1 = a_1 a_2 \dots a_n$ be a permutation of $[n]$ and let $\pi_2 = b_1 b_2 \dots b_k$ be a permutation of $[k]$. We say that π_1 contains a type π_2 subsequence if there exists $1 \leq i_{b_1} < i_{b_2} < \dots < i_{b_k} \leq n$ such that $a_{i_1} < a_{i_2} < \dots < a_{i_k}$. We say π_1 avoids π_2 if π_1 contains no π_2 -subsequence. For example, a permutation π avoids 231 if it cannot be written as $\dots a \dots b \dots c \dots$ so that $c < a < b$.

Tarjan [24] proved that a permutation is 1-stack-sortable if and only if it avoids the pattern 231. West [26, 27] proved that a permutation fails to be 2-stack-sortable if it contains a subsequence of type 2341 or a subsequence of type 3241 which is not

part of a subsequence of type 35241 and it is two-stack-sortable if it contains no such subsequence. West also conjectured the number of 2-stack-sortable permutations of $[n]$ to be [27, 26]

$$\frac{2(3n)!}{(n+1)!(2n+1)!}.$$

The conjecture was first proved by D. Zeilberger [28]. Later, two bijections between two-stack-sortable permutations and non-separable planar graphs were given by S. Dulucq, S. Gire and O. Guibert [7]; I. P. Goulden and J. West [11]. More contributions to this problem were given by Miklós Bóna, Mireille Bousquet-Mélou, Leopold Travis and others [2, 3, 4, 5, 6, 25].

1.1. The generalization

In this thesis, we generalize the ordinary permutations to r -permutations [13, 14, 16, 17, 18] and then enumerate the number of 2-stack-sortable r -permutations under the generalized stack-sorting operation on r -permutations.

DEFINITION 1.1.1. If a permutation $a_1 a_2 \cdots a_{rn}$ of $\{1^r, 2^r, \dots, n^r\}$ satisfies the condition that if $i < j < k$ and $a_i = a_k$ then $a_j \leq a_i$, we call it an r -permutation of $[n]$.

DEFINITION 1.1.2. Let π be an r -permutation of $[n]$. For $n > 0$, we can write π as $\pi = \alpha_1 n \alpha_2 n \cdots n \alpha_{r+1}$. The stack-sorting operation S on r -permutations is defined by

$$S(\pi) = S(\alpha_1) S(\alpha_2) \cdots S(\alpha_{r+1}) n.$$

When $n = 0$, i.e., when π is an empty permutation, we define $S(\pi)$ to be $\emptyset$. Note that $S(\pi)$ is an ordinary permutation.

DEFINITION 1.1.3. Given an r -permutation π and a letter a in π , we call a a type i descent ($i = 1, \dots, r$) if the i th occurrence of a is immediately followed by a smaller letter, and we call a a type 0 descent if the first occurrence of a immediately follows a smaller letter. We denote the set of type i descents of an r -permutation π by $\pi^{(i)}$ ($i = 0, \dots, r$).

REMARK 1.1.4. Notice that the r -permutations are usually defined the other way, i.e., a permutation $a_1 a_2 \cdots a_{rn}$ of $\{1^r, 2^r, \dots, n^r\}$ is an r -permutation if it satisfies the condition that if $i < j < k$ and $a_i = a_k$, then $a_j \geq a_i$. The reason for the change is that we expect $S(\pi)$ to be an ordinary permutation for an r -permutation π , while the traditional definition of r -permutation can not give us that. For example, 123321 is a 2-permutation under the traditional definition, but $S(123321) = 121233$. This behavior is very different from what we study here.

REMARK 1.1.5. Note that a type 0 descent is actually an ascent.

REMARK 1.1.6. Often, the last letter (or, the last position) of a permutation is considered to be a descent but not here.

EXAMPLE 1.1.7. When $r = 3$, $n = 6$,

$$\pi = 544453222335611166$$


is a 3-permutation of $\{1, 2, \dots, 6\}$. Then

$$\begin{aligned}
 S(\pi) &= S(544453222335)S(111)6 \\
 &= S(444)S(322233)516 \\
 &= 4S(222)3516 \\
 &= 423516.
 \end{aligned}$$

The sets of type i descents ($i = 0, \dots, 3$) of π are $\pi^{(0)} = \{6\}$, $\pi^{(1)} = \{3, 5, 6\}$, $\pi^{(2)} = \{5\}$ and $\pi^{(3)} = \emptyset$.

The r -permutations can be represented as $(r + 1)$ -ary decreasing trees. If we have an r -permutation π of $[n]$ and $\pi = \alpha_1 n \alpha_2 n \cdots n \alpha_{r+1}$, to get the tree representation of π , we set n to be the root of the tree and recursively set the tree representation of α_i ($i = 1, 2, \dots, r + 1$) to be the i th child of the root. If α_i is nonempty, then π has an i th child and n is a type $i - 1$ descent. Therefore it is clear that each i th child corresponds to a type $i - 1$ descent. For example, the 3-permutation above can be represented as in Figure 1.1.

There are three kinds of traversals of a tree: preorder, inorder, and postorder [23, p. 243; 15, p. 315]. In a preorder traversal of a tree, the root is visited first and then the subtrees rooted at its children are traversed recursively. A postorder traversal recursively traverses the subtrees rooted at the children of the root first, and then visits the root. An inorder traversal, in our case, recursively traverses the subtree rooted at the first child of the root, then visits the root, then recursively traverses the subtree rooted at the second child of the root, then visits the root, and so on. Hence, if we read the tree in Figure 1.1 in inorder, we get back the original r -permutation π . If we read it in postorder, we get an ordinary permutation and it corresponds to $S(\pi)$.

FIGURE 1.1. Tree representation of π

Because any $(r + 1)$ -ary tree on n vertices has $n - 1$ children, we have

LEMMA 1.1.8. *Every r -permutation of $[n]$ has a total of $n - 1$ descents of types $0, 1, \dots, r$.*

1.2. Stack-sortable r -permutations

Now let us first consider the enumeration of 1-stack-sortable r -permutations (which we call here stack-sortable r -permutations) with all types of descents.

We weight any type i descent by x_i . Therefore the weight of the r -permutation π in Example 1.1.7 is $x_0 x_1^3 x_2$. Because of Lemma 1.1.8, we do not need a parameter to keep track of the number of different letters in an r -permutation.

For any unlabeled $(r + 1)$ -ary tree, there is one and only one way to label the nodes to make it an $(r + 1)$ -ary decreasing tree such that if we read it in postorder, we

get an identity ordinary permutation. That gives a bijection between stack-sortable r -permutations and unlabeled $(r+1)$ -ary trees. Therefore, if we weight an unlabeled $(r+1)$ -ary tree by weighting any i th child by x_{i-1} , then the total weight of stack-sortable r -permutations is the total weight of unlabeled $(r+1)$ -ary trees. Let $A(x)$ be the weight of all stack-sortable r -permutations. Then we have

$$A(x) = \prod_{i=0}^r (1 + x_i A(x))$$

where $x = (x_0, x_1, \dots, x_r)$.

To solve this functional equation, we use Lagrange inversion [10, p. 21; 9].

LEMMA 1.2.1. (*Multivariable Lagrange Inversion*) Let $f(\lambda) \in \mathbb{R}[[\lambda]]$ and $\phi_1(\lambda), \dots, \phi_m(\lambda) \in \mathbb{R}[[\lambda]]$ where $\lambda = (\lambda_1, \dots, \lambda_m)$. Suppose that $w_i = t_i \phi_i(w)$ for $i = 1, \dots, m$, where $w = (w_1, \dots, w_m)$. Let $\phi = (\phi_1, \dots, \phi_m)$, $t = (t_1, \dots, t_m)$ and $k = (k_1, \dots, k_m)$. Then

$$f(w(t)) = \sum_k t^k [\lambda^k] \left\{ f(\lambda) \phi^k(\lambda) \left| \delta_{ij} - \frac{\lambda_j}{\phi_i(\lambda)} \frac{\partial \phi_i(\lambda)}{\partial \lambda_j} \right| \right\},$$

where $[\lambda^k]f(\lambda)$ is the coefficient of $\lambda_1^{k_1} \cdots \lambda_m^{k_m}$ of the formal power series $f(\lambda)$.

In our case, we let $w_i = x_i A(x)$ and $\phi_i(\lambda) = \prod_{j=0}^r (1 + \lambda_j)$ for $i = 0, \dots, r$. Then w_i satisfy the condition that $w_i = x_i \phi_i(w)$. Also let $f(\lambda) = \prod_{j=0}^r (1 + \lambda_j)$. Then $f(w) = A(x)$. Using multivariable Lagrange inversion, we get

$$\begin{aligned} A(x) &= f(w) \\ &= \sum_k x^k [\lambda^k] \left\{ \left(\prod_{i=0}^r (1 + \lambda_i) \right)^{k_0 + \dots + k_r + 1} \left| \delta_{ij} - \frac{\lambda_j}{1 + \lambda_j} \right| \right\}. \end{aligned}$$

Now we introduce some notation. Let

$$(1.2.1) \quad e_k(x) = \sum_{0 \leq i_1 < i_2 < \dots < i_k \leq r} x_{i_1} x_{i_2} \cdots x_{i_k}$$

be the k th elementary symmetric function of $x = (x_0, x_1, \dots, x_r)$. Define

$$\begin{aligned}
 (1.2.2) \quad E(x, u) &= E(x_0, x_1, \dots, x_r; u) \\
 &= \prod_{i=0}^r (1 + x_i u) \\
 &= \sum_{i=0}^r e_i(x) u^i
 \end{aligned}$$

and let

$$(1.2.3) \quad E(x) = E(x, 1).$$

Also we introduce the following lemma to evaluate the determinant.

LEMMA 1.2.2. *Let $M(x) = |\delta_{ij} + (\delta_{ij} - 1)x_j|$, where $0 \leq i, j \leq r$, be the determinant of the $(r+1) \times (r+1)$ matrix. Then*

$$M(x) = E(x) \left(1 - \sum_{i=0}^r \frac{x_i}{1 + x_i} \right).$$

PROOF.

$$M(x) = \begin{vmatrix}
 1 & -x_1 & -x_2 & \cdots & -x_{r-1} & -x_r \\
 -x_0 & 1 & -x_2 & \cdots & -x_{r-1} & -x_r \\
 \dots\dots\dots & & & & & \\
 -x_0 & -x_1 & -x_2 & \cdots & 1 & -x_r \\
 -x_0 & -x_1 & -x_2 & \cdots & -x_{r-1} & 1
 \end{vmatrix}.$$

Subtracting the last row from every other row, we get

$$M(x) = \begin{vmatrix} 1+x_0 & 0 & 0 & \cdots & 0 & -(1+x_r) \\ 0 & 1+x_1 & 0 & \cdots & 0 & -(1+x_r) \\ 0 & 0 & 1+x_2 & \cdots & 0 & -(1+x_r) \\ \dots\dots\dots & & & & & \\ 0 & 0 & 0 & \cdots & 1+x_{r-1} & -(1+x_r) \\ -x_0 & -x_1 & -x_2 & \cdots & -x_{r-1} & 1 \end{vmatrix}.$$

If we let $x' = (x_1, x_2, \dots, x_r)$, then by expanding the first column, we get

$$\begin{aligned} M(x) &= (1+x_0) \begin{vmatrix} 1+x_1 & 0 & \cdots & -(1+x_r) \\ 0 & 1+x_2 & \cdots & -(1+x_r) \\ \dots\dots\dots & & & \\ -x_1 & -x_2 & \cdots & 1 \end{vmatrix} \\ &\quad - (-1)^r x_0 \begin{vmatrix} 0 & 0 & \cdots & 0 & -(1+x_r) \\ 1+x_1 & 0 & \cdots & 0 & -(1+x_r) \\ \dots\dots\dots & & & & \\ 0 & 0 & \cdots & 1+x_{r-1} & 1 \end{vmatrix} \\ &= (1+x_0)M(x') - (-1)^r x_0 (-1)^r \prod_{i=1}^r (1+x_i) \\ &= (1+x_0)M(x') - x_0 \prod_{i=1}^r (1+x_i). \end{aligned}$$

Now we use induction. It is easy to see that the lemma is true when $r = 0$. Suppose the lemma is true for $M(x')$, i.e.,

$$M(x') = E(x') \left(1 - \sum_{i=1}^r \frac{x_i}{1+x_i} \right).$$

Then

$$\begin{aligned}
M(x) &= (1 + x_0)E(x') \left(1 - \sum_{i=1}^r \frac{x_i}{1 + x_i} \right) - x_0 \prod_{i=1}^r (1 + x_i) \\
&= E(x) \left(1 - \sum_{i=1}^r \frac{x_i}{1 + x_i} \right) - E(x) \frac{x_0}{1 + x_0} \\
&= E(x) \left(1 - \sum_{i=0}^r \frac{x_i}{1 + x_i} \right).
\end{aligned}$$

The lemma is proved. □

With Lemma 1.2.2, it follows that

$$\begin{aligned}
A(x) &= \sum_k x^k [\lambda^k] \left\{ \left(\prod_{i=0}^r (1 + \lambda_i) \right)^{k_0 + \dots + k_r + 1} \left(\sum_{i=0}^r \frac{1}{1 + \lambda_i} - r \right) \right\} \\
&= \sum_k x^k \left\{ \sum_i \binom{n}{k_1} \dots \binom{n-1}{k_i} \dots \binom{n}{k_r} - r \prod_i \binom{n}{k_i} \right\} \\
&= \sum_k x^k \left\{ \left(\sum_i \frac{n - k_i}{n} - r \right) \prod_i \binom{n}{k_i} \right\} \\
&= \sum_k x^k \frac{1}{n} \prod_{i=0}^r \binom{n}{k_i}
\end{aligned}$$

where $n = 1 + k_0 + \dots + k_r$. Therefore,

THEOREM 1.2.3. *The number of stack-sortable r -permutations with k_i descents of type i is*

$$\frac{1}{n} \prod_{i=0}^r \binom{n}{k_i}$$

where $n = 1 + k_0 + \dots + k_r$.

REMARK 1.2.4. This functional equation can also be solved by one variable Lagrange inversion by introducing a new variable, i.e., letting

$$A(z) = z \prod_{i=0}^r (1 + x_i A(z)).$$

REMARK 1.2.5. These numbers also come up in counting noncrossing partitions [21, 20, 8].

REMARK 1.2.6. When $r = 1$, these numbers are Narayana numbers [22, 12].

CHAPTER 2

Two-stack-sortable r -permutations

2.1. The functional equations

Now let us consider the case of two-stack-sortable r -permutations. An r -permutation π of $[n]$ is two-stack-sortable if $S^2(\pi)$ is an identity permutation. Let $\pi = \alpha_1 n \alpha_2 n \cdots n \alpha_{r+1}$ where n is the largest element in π . Then

$$S(\pi) = S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{r+1})n$$


$$S^2(\pi) = S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{r+1}))n$$

Denote the identity permutation by I . (Here we abuse the notation by letting I be the identity permutation of $[n]$ for any positive integer n .) One can notice that if $S^2(\pi) = I$ and if m is the largest number in $\alpha_1 \alpha_2 \cdots \alpha_{r+1}$, then m can occur in only one of the α_i because if m appeared in two of the α_i , the r -permutation condition for π would be violated. Thus $(\alpha_1, \alpha_2, \dots, \alpha_{r+1})$ is an $(r+1)$ -tuples of r -permutations satisfying the condition that

$$S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{r+1})) = I.$$

This suggests that we study the subject defined as follows.

DEFINITION 2.1.1. $(\alpha_1, \alpha_2, \dots, \alpha_k)$ is a k -tuple r -permutation of $[n]$ if $\alpha_1 \alpha_2 \cdots \alpha_k$ is an r -permutation of $[n]$ and any letter appears in only one of the α_i . Also we call α_i the i th component of the k -tuple r -permutation.

FIGURE 2.1. Forest representation of π

We call a k -tuple r -permutation $(\alpha_1, \alpha_2, \dots, \alpha_k)$ a two-stack-sortable k -tuple r -permutation if it satisfies the condition that

$$S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_k)) = I.$$

From the definition, we know that each component α_i of a k -tuple r -permutation is itself an r -permutation so that we can interpret a k -tuple r -permutation of $[n]$ as a forest of k $(r+1)$ -ary decreasing trees. The set of the descents of type i of k -tuple r -permutations $\pi = (\alpha_1, \dots, \alpha_k)$ is the union of the sets of descents of type i of α_j ($j = 1, \dots, k$), i.e.

$$\pi^{(i)} = \bigcup_{j=1}^k \alpha_j^{(i)}.$$

For example, let $\pi = (\alpha_1, \alpha_2)$ where $\alpha_1 = 553111335$ and $\alpha_2 = 442224776667$. Then (α_1, α_2) is a 2-tuple 3-permutation of $\{1, 2, \dots, 7\}$ and it can be represented as in Figure 2.1. Since

$$\begin{aligned} S(S(\alpha_1)S(\alpha_2)) &= S(1352467) \\ &= 1324567, \end{aligned}$$

π is not two-stack-sortable. The sets of descents of π are $\pi^{(0)} = \{7\}$, $\pi^{(1)} = \{3\}$, $\pi^{(2)} = \{4, 5, 7\}$, $\pi^{(3)} = \emptyset$.

Let $g_n^{(k)}$ be the sum of the weights of two-stack-sortable k -tuple r -permutations of n different letters *such that every component is nonempty*. We need every component to be nonempty so that we can keep track of every type of descent. Suppose $(\alpha_1, \alpha_2, \dots, \alpha_k)$ is a two-stack-sortable k -tuple r -permutation of $[n]$ in which every component is nonempty and the largest element n appears in α_l . Let $\alpha_l = \beta_1 n \beta_2 n \cdots n \beta_{r+1}$. Then

(2.1.1)

$$\begin{aligned} I &= S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_k)) \\ &= S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{l-1})S(\beta_1)S(\beta_2) \cdots S(\beta_{r+1}) n S(\alpha_{l+1}) \cdots S(\alpha_k)) \\ &= S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{l-1})S(\beta_1)S(\beta_2) \cdots S(\beta_{r+1})) S(S(\alpha_{l+1}) \cdots S(\alpha_k)) n. \end{aligned}$$

Therefore, $(\alpha_1, \dots, \alpha_{l-1}, \beta_1, \dots, \beta_{r+1})$ and $(\alpha_{l+1}, \dots, \alpha_k)$ are both two-stack-sortable tuples of r -permutations.

Let us first consider the case when $l = 1$. By (2.1.1), we have

$$I = S(S(\beta_1)S(\beta_2) \cdots S(\beta_{r+1})) S(S(\alpha_2) \cdots S(\alpha_k)) n.$$

Suppose there are $n - i$ letters (including n) in α_1 . Then the weight of $(\alpha_2, \alpha_3, \dots, \alpha_k)$ is a term of $g_i^{(k-1)}$. For $(\beta_1, \dots, \beta_{r+1})$, if all β_j are nonempty, then the weight of $(\beta_1, \dots, \beta_{r+1})$ is a term of $g_{n-1-i}^{(r+1)}$. But by the definition of the descents of r -permutations, there is a descent of each type in α_1 for every appearance of n . So the weight of α_1 is $x_0 x_1 \cdots x_r$ times the weight of $(\beta_1, \dots, \beta_{r+1})$. Therefore the sum of the weights of all such α_1 is $e_{r+1}(x) g_{n-1-i}^{(r+1)}$. If only one of the β_j is empty, then the weight of $(\beta_1, \dots, \beta_{r+1})$ is a term of $g_{n-1-i}^{(r)}$. Again, by the definition of the descents of r -permutations, there

are $r - 1$ descents of different types for $r - 1$ appearances of n . So, if β_{j_0} is empty but β_j is not empty for $j \neq j_0$, then the weight of α_1 is $x_0 x_1 \cdots \hat{x}_{j_0} \cdots x_r$ times the weight of $(\beta_1, \dots, \beta_{r+1})$. Therefore the sum of the weights of all such α_1 is $e_r(x) g_{n-1-i}^{(r)}$. In general, if there are m of the β_j that are nonempty, then the sum of the weights of all such α_1 is $e_m(x) g_{n-1-i}^{(m)}$. Therefore, the sum of weights of all the k -tuple r -permutation $(\alpha_1, \alpha_2, \dots, \alpha_k)$ in which n is in α_1 is

$$\sum_{i=0}^{n-1} g_i^{(k-1)} [g_{n-1-i}^{(0)} + g_{n-1-i}^{(1)} e_1(x) + \cdots + g_{n-1-i}^{(r+1)} e_{r+1}(x)],$$

For the same reason, when n is in α_2 , we have

$$I = S(S(\alpha_1)S(\beta_1)S(\beta_2) \cdots S(\beta_{r+1})) S(S(\alpha_3) \cdots S(\alpha_k)) n.$$

Suppose there are $n - i$ letters in (α_1, α_2) . Then the sum of the weights of all the k -tuple r -permutation $(\alpha_1, \alpha_2, \dots, \alpha_k)$ in which n is in α_2 is

$$\sum_{i=0}^{n-1} g_i^{(k-2)} [g_{n-1-i}^{(1)} + g_{n-1-i}^{(2)} e_1(x) + \cdots + g_{n-1-i}^{(r+2)} e_{r+1}(x)],$$

and so on. Thus we have the crucial recursive formula

$$(2.1.2) \quad g_n^{(k)} = \sum_{i=0}^{n-1} \sum_{l=0}^{k-1} \sum_{j=0}^{r+1} g_{n-1-i}^{(l+j)} g_i^{(k-1-l)} e_j(x)$$

with the initial conditions that $g_0^{(k)} = \delta_{0k}$ and $g_n^{(0)} = \delta_{n0}$.

Now let

$$(2.1.3) \quad f_n^{(k)} = \sum_{j=0}^{r+1} g_n^{(k+j)} e_j(x).$$

Then

$$(2.1.4) \quad g_n^{(k)} = f_{n-1}^{(k-1)} + \sum_{i=0}^{n-1} f_{n-1-i}^{(k-2)} g_i^{(1)} + \cdots + \sum_{i=0}^{n-1} f_{n-1-i}^{(0)} g_i^{(k-1)}.$$

Following the analysis above one can easily see that $f_n^{(k)}$ is the sum of the weights of $(k+1)$ -tuple r -permutations $(\alpha_1, \alpha_2, \dots, \alpha_{k+1})$ of $[n+1]$ such that the α_i are all nonempty, $S(S(\alpha_1)S(\alpha_2) \cdots S(\alpha_{k+1})) = I$ and the largest letter only appears in α_{k+1} .

Also let

$$(2.1.5) \quad f_k = \sum_{n=0}^{\infty} f_n^{(k)}, \quad F = \sum_{k=0}^{\infty} f_k z^k,$$

$$(2.1.6) \quad g_k = \sum_{n=0}^{\infty} g_n^{(k)}, \quad G = \sum_{k=0}^{\infty} g_k z^k.$$

Then it follows that

$$g_0 = 1,$$

$$f_0 = g_1,$$

$$(2.1.7) \quad f_k = g_k + g_{k+1}e_1(x) + \cdots + g_{r+k+1}e_{r+1}(x),$$

$$(2.1.8) \quad g_k = f_{k-1}g_0 + f_{k-2}g_1 + \cdots + f_0g_{k-1},$$

and from (2.1.8), we get

$$(2.1.9) \quad G = 1 + zFG.$$

Notice that if we let r go to infinity, we can interpret k -tuple r -permutations in terms of forests of decreasing trees and equations (2.1.7) to (2.1.9) are still well defined. Therefore, we have the functional equations as follows.

THEOREM 2.1.2. *Let F and G be defined as in (2.1.5) and (2.1.6). Then F and G satisfy the functional equations*

$$(2.1.10) \quad f_k = \sum_{j=0}^{\infty} g_{k+j} e_j(x) \text{ for } k \geq 0,$$

$$(2.1.11) \quad G = 1 + zFG$$

and F and G are uniquely determined as power series by these functional equations.

REMARK 2.1.3. From (2.1.4), we see that $g_n^{(k)}$ are uniquely determined by $f_i^{(j)}$ and $g_i^{(j)}$ where $i < n$. From (2.1.3), we see that $f_n^{(k)}$ are uniquely determined by $g_n^{(j)}$. Therefore, given $g_0^{(0)} = 1$ and $g_0^{(k)} = 0$ when $k \geq 1$, all $g_n^{(k)}$ and $f_n^{(k)}$ are uniquely determined by (2.1.3) and (2.1.4). Hence, F and G are uniquely determined by the functional equations in Theorem 2.1.2.

REMARK 2.1.4. Notice that $g_1 = f_0$ is the generating function of two-stack-sortable r -permutations.

REMARK 2.1.5. Notice that in the ring $\mathbb{Q}[[z, \frac{x_0}{z}, \frac{x_1}{z}, \frac{x_2}{z}, \dots]]$,

$$G \cdot E\left(x, \frac{1}{z}\right) = \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} g_{k+j} e_j(x) z^k.$$

Therefore, by (2.1.10), $F - G \cdot E\left(x, \frac{1}{z}\right)$ has only negative powers of z . This is the key to solving the functional equations (2.1.10) and (2.1.11).

2.2. The solution to the functional equations

Let $y = (y_0, y_1, y_2, \dots)$, where the y_i are uniquely determined as formal power series in the x_i by

$$x_j = \frac{y_j(1 + y_j)}{\prod_{i=0}^{\infty} (1 + y_i)^2} = \frac{y_j(1 + y_j)}{E^2(y)}.$$

Let $t = zE^2(y)$ where $E(y)$ is defined as in (1.2.2) and (1.2.3). Let $c(t) = \frac{1 - \sqrt{1 - 4t}}{2t}$ be the generating function of the Catalan numbers. Then

THEOREM 2.2.1. *The solution to the functional equations in Theorem 2.1.2 is:*

$$(2.2.1) \quad G = \frac{c(t)E(y)}{E(y, c(t))},$$

$$(2.2.2) \quad F = \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right).$$

The proof of the theorem consists of the following lemmas.

LEMMA 2.2.2. *In the ring $\mathbb{Q}[[z, \frac{x_0}{z}, \frac{x_1}{z}, \dots]]$,*

$$E\left(x, \frac{1}{z}\right) = E(y, c(t))E\left(y, \frac{1}{tc(t)}\right)$$

PROOF. First, we notice that $E\left(x, \frac{1}{z}\right) \in \mathbb{Q}[[z, \frac{x_0}{z}, \frac{x_1}{z}, \dots]]$ and $E\left(y, \frac{1}{tc(t)}\right) \in \mathbb{Q}[[t, \frac{y_0}{t}, \frac{y_1}{t}, \dots]]$. It is also easy to verify that $\mathbb{Q}[[z, \frac{x_0}{z}, \frac{x_1}{z}, \dots]] = \mathbb{Q}[[t, \frac{y_0}{t}, \frac{y_1}{t}, \dots]]$. Therefore, since $c(t)$ satisfies $tc^2(t) - c(t) + 1 = 0$,

$$\begin{aligned} (1 + c(t)y_i) \left(1 + \frac{y_i}{tc(t)} \right) &= 1 + c(t)y_i + \frac{y_i}{tc(t)} + \frac{y_i^2}{t} \\ &= 1 + \frac{y_i + y_i^2}{t} \\ &= 1 + \frac{x_i}{z}. \end{aligned}$$

Hence,

$$\begin{aligned}
E\left(x, \frac{1}{z}\right) &= \prod_{i=0}^{\infty} \left(1 + \frac{x_i}{z}\right) \\
&= \prod_{i=0}^{\infty} \left(1 + \frac{y_i + y_i^2}{t}\right) \\
&= \prod_{i=0}^{\infty} (1 + c(t)y_i) \left(1 + \frac{y_i}{tc(t)}\right) \\
&= E(y, c(t)) E\left(y, \frac{1}{tc(t)}\right).
\end{aligned}$$

□

LEMMA 2.2.3. *Let*

$$\begin{aligned}
\hat{G} &= \frac{c(t)E(y)}{E(y, c(t))}, \\
\hat{F} &= \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right).
\end{aligned}$$

Then $\hat{G}$ and $\hat{F}$ satisfy

$$\hat{G} = 1 + z\hat{F}\hat{G}.$$

PROOF.

$$\begin{aligned}
1 - z\hat{F} &= 1 - z \cdot \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right) \\
&= 1 - \frac{zE^2(y)}{t} \left(1 - \frac{E(y, c(t))}{c(t)E(y)} \right) \\
&= \frac{E(y, c(t))}{c(t)E(y)} \\
&= \frac{1}{\hat{G}}
\end{aligned}$$

□

LEMMA 2.2.4. *In the ring $\mathbb{Q}[[z, \frac{x_0}{z}, \frac{x_1}{z}, \frac{x_2}{z}, \dots]]$, define*

$$\hat{F}^- = \hat{G} \cdot E\left(x, \frac{1}{z}\right) - \hat{F}.$$

Then $\hat{F}^-$ has only negative powers of z .

PROOF.

$$\begin{aligned} \hat{F}^- &= \hat{G} \cdot E\left(x, \frac{1}{z}\right) - \hat{F} \\ &= \frac{c(t)E(y)}{E(y, c(t))} \cdot E\left(x, \frac{1}{z}\right) - \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)}\right) \\ &= \frac{c(t)E(y)}{E(y, c(t))} \cdot E(y, c(t))E\left(y, \frac{1}{tc(t)}\right) - \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)}\right) \\ &= \frac{E(y)}{t} \left[tc(t)E\left(y, \frac{1}{tc(t)}\right) + \frac{E(y, c(t))}{c(t)} - E(y)\right] \\ &= \frac{E(y)}{t} \sum_{n=0}^{\infty} [c(t)^{n-1} + t^{1-n}c(t)^{1-n} - 1] e_n(y) \\ &= E(y) \sum_{n=0}^{\infty} \left[\frac{1}{t^n} [(tc(t))^{n-1} + c(t)^{1-n}] - \frac{1}{t}\right] e_n(y). \end{aligned}$$

Since

$$\begin{aligned} (2.2.3) \quad (tc(t))^{n-1} + c(t)^{1-n} &= \left(\frac{1 - \sqrt{1-4t}}{2}\right)^{n-1} + \left(\frac{1 + \sqrt{1-4t}}{2}\right)^{n-1} \\ &= \frac{1}{2^{n-2}} \sum_{k \text{ even}} (\sqrt{1-4t})^k \binom{n-1}{k} \\ &= \frac{1}{2^{n-2}} \sum_{k \text{ even}} (1-4t)^{k/2} \binom{n-1}{k} \end{aligned}$$

is a polynomial of degree less than or equal to $(n-1)/2$, $\hat{F}^-$ has only negative powers of t and thus has only negative powers of z . $\square$

LEMMA 2.2.5. *Let $\hat{G}$ and $\hat{F}$ be defined as above. Define*

$$\hat{f}_k = [z^k]\hat{F}, \quad \hat{g}_k = [z^k]\hat{G}.$$

Then $\hat{g}_k$ and $\hat{f}_k$ satisfy the functional equation that

$$\hat{f}_k = \sum_{j=0}^{\infty} \hat{g}_{k+j} e_j(x) \text{ for } k \geq 0.$$

PROOF. From the definition of $\hat{f}_k$ and $\hat{g}_k$, we know that

$$\begin{aligned} \hat{F} &= \sum_{k=0}^{\infty} \hat{f}_k z^k, \\ \hat{G} &= \sum_{k=0}^{\infty} \hat{g}_k z^k. \end{aligned}$$

Then

$$[z^k] \left\{ \hat{G} \cdot E \left(x, \frac{1}{z} \right) - \hat{F} \right\} = \sum_{j=0}^{\infty} \hat{g}_{k+j} e_j(x) - \hat{f}_k.$$

By Lemma 2.2.4, this sum is zero when $k \geq 0$.

The lemma is proved. □

Now Theorem 2.2.1 follows from Lemma 2.2.3 and Lemma 2.2.5.

2.3. The number of two-stack-sortable r -permutations with descents

We know that the generating function for two-stack-sortable r -permutations is

$$\begin{aligned} g_1 &= f_0 \\ &= [z^0] F \\ &= [t^0] \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right) \\ &= [t^0] \left\{ E(y) \sum_{n=0}^{\infty} \frac{1}{t} (1 - c^{n-1}(t)) e_n(y) \right\} \end{aligned}$$

$$\begin{aligned}
&= [t^0] \left\{ E(y) \sum_{n=0}^{\infty} \sum_{i=0}^{\infty} \frac{1-n}{i+1} \binom{n+2i}{i} e_n(y) t^i \right\} \\
&= E(y) \sum_{n=0}^{\infty} (1-n) e_n(y)
\end{aligned}$$

where

$$x_i = \frac{y_i(1+y_i)}{E^2(y)}.$$

We can now evaluate the number of two-stack-sortable r -permutations with the help of multivariable Lagrange inversion.

THEOREM 2.3.1. *The number of two-stack-sortable r -permutations with k_i descents of type i is*

$$(2.3.1) \quad \frac{1}{n^2} \prod_{i=0}^r \frac{n}{n-k_i} \binom{2n-1-k_i}{k_i},$$

where $n = 1 + k_0 + k_1 + \cdots + k_r$.

First, we introduce the following identity:

LEMMA 2.3.2.

$$\sum_{i=0}^{r+1} (1-i) e_i(y) = E(y) \left(1 - \sum_{i=0}^r \frac{y_i}{1+y_i} \right).$$

PROOF. Since $e_i(y)$ is a homogeneous polynomial in $y_0, y_1, \dots, y_r$ of degree i ,

$$i e_i(y) = \sum_{j=0}^r y_j \frac{\partial e_i(y)}{\partial y_j}.$$

Also notice that

$$\frac{\partial E(y)}{\partial y_j} = \frac{E(y)}{1+y_j}.$$

Thus

$$\begin{aligned}
\sum_{i=0}^{r+1} (1-i)e_i(y) &= E(y) - \sum_{i=0}^{r+1} ie_i(y) \\
&= E(y) - \sum_{i=0}^{r+1} \sum_{j=0}^r \frac{y_j \partial e_i(y)}{\partial y_j} \\
&= E(y) - \sum_{j=0}^r y_j \sum_{i=0}^{r+1} \frac{\partial e_i(y)}{\partial y_j} \\
&= E(y) - \sum_{j=0}^r y_j \frac{E(y)}{1+y_j} \\
&= E(y) \left(1 - \sum_{j=0}^r \frac{y_j}{1+y_j} \right).
\end{aligned}$$

□

Therefore,

$$\begin{aligned}
(2.3.2) \quad g_1 &= E(y) \sum_{n=0}^{\infty} (1-n)e_n(y) \\
&= E^2(y) \left(1 - \sum_{j=0}^r \frac{y_j}{1+y_j} \right).
\end{aligned}$$

If for any $y = (y_0, y_1, \dots, y_r)$, we define

$$\begin{aligned}
A(y, u) &= A(y_0, \dots, y_r; u) \\
&= 1 - \sum_{i=0}^r \frac{y_i u}{1+y_i u},
\end{aligned}$$

and let

$$A(y) = A(y, 1),$$

then $g_1 = E^2(y)A(y)$.

The proof of Theorem 2.3.1 now consists of following lemmas.

Notice that for any formal power series $f(x) = \sum_k a_k x^k \in \mathbb{Q}[[x_0, x_1, \dots, x_r]]$ (where $k = (k_0, \dots, k_r)$ and $x = (x_0, \dots, x_r)$), if we define operators $D_x, I_x: \mathbb{Q}[[x_0, x_1, \dots, x_r]] \rightarrow \mathbb{Q}[[x_0, x_1, \dots, x_r]]$ by

$$D_x(f(x)) = \sum_{i=0}^r x_i \frac{\partial f(x)}{\partial x_i},$$

$$I_x(f(x)) = f(x),$$

then $(D_x + I_x)(f(x)) = \sum_k n a_k x^k$, where $n = k_0 + \dots + k_r + 1$. Therefore, to prove

$$[x^k]g_1 = \frac{1}{n^2} \prod_{i=0}^r \frac{n}{n - k_i} \binom{2n - 1 - k_i}{k_i},$$

we only need to prove that

$$[x^k](D_x + I_x)^2 g_1 = \prod_{i=0}^r \frac{n}{n - k_i} \binom{2n - 1 - k_i}{k_i}.$$

Let $u_i = \sum_{j=0}^r \frac{x_j \partial y_i}{\partial x_j}$. Then

LEMMA 2.3.3.

$$u_i = \frac{y_i(1 + y_i)}{1 + 2y_i} \frac{1}{A(y, 2)}$$

PROOF. Take the logarithm of both sides of $x_i = \frac{y_i(1 + y_i)}{E^2(y)}$. We get

$$(2.3.3) \quad \ln x_i = \ln y_i + \ln(1 + y_i) - 2 \ln E(y).$$

Then by differentiating both sides of 2.3.3 with respect to x_j , we get

$$(2.3.4) \quad \frac{1}{x_i} = \frac{1 + 2y_i}{y_i(1 + y_i)} \frac{\partial y_i}{\partial x_i} - \frac{2}{E(y)} \frac{\partial E(y)}{\partial x_i}, \quad \text{if } j = i,$$

$$(2.3.5) \quad 0 = \frac{1 + 2y_i}{y_i(1 + y_i)} \frac{\partial y_i}{\partial x_j} - \frac{2}{E(y)} \frac{\partial E(y)}{\partial x_j}, \quad \text{if } j \neq i.$$

Now multiplying x_j by both (2.3.4) and (2.3.5) and summing on j , we get

$$\begin{aligned}
 1 &= \frac{1+2y_i}{y_i(1+y_i)} \sum_{j=0}^r \frac{x_j \partial y_i}{\partial x_j} - \frac{2}{E(y)} \sum_{j=0}^r \frac{x_j \partial E(y)}{\partial x_j} \\
 (2.3.6) \quad &= \frac{1+2y_i}{y_i(1+y_i)} u_i - \frac{2}{E(y)} D_x(E(y)).
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 D_x(E(y)) &= \sum_{j=0}^r x_j \frac{\partial E(y)}{\partial x_j} \\
 &= \sum_{j=0}^r x_j \sum_{i=0}^r \frac{\partial E(y)}{\partial y_i} \frac{\partial y_i}{\partial x_j} \\
 &= E(y) \sum_{i=0}^r \frac{1}{1+y_i} \sum_{j=0}^r x_j \frac{\partial y_i}{\partial x_j} \\
 (2.3.7) \quad &= E(y) \sum_{i=0}^r \frac{1}{1+y_i} u_i
 \end{aligned}$$

By solving (2.3.6) and (2.3.7) for u_i , we get

$$u_i = \frac{y_i(1+y_i)}{1+2y_i} \frac{1}{A(y, 2)}.$$

The lemma is proved. □

Using the lemma above, the proofs of the following lemmas are just a matter of simple computations.

LEMMA 2.3.4.

$$D_x(E^2(y)) = \frac{E^2(y)}{A(y, 2)} - E^2(y).$$

PROOF.

$$D_x(E^2(y)) = \sum_{j=0}^r x_j \frac{\partial E^2(y)}{\partial x_j}$$

$$\begin{aligned}
&= \sum_{j=0}^r x_j \sum_{i=0}^r \frac{\partial E^2(y)}{\partial y_i} \frac{\partial y_i}{\partial x_j} \\
&= \sum_{i=0}^r \frac{\partial E^2(y)}{\partial y_i} u_i \\
&= \sum_{i=0}^r \frac{2E^2(y)}{1+y_i} \frac{y_i(1+y_i)}{1+2y_i} \frac{1}{A(y, 2)} \\
&= \frac{E^2(y)}{A(y, 2)} \sum_{i=0}^r \frac{2y_i}{1+2y_i} \\
&= \frac{E^2(y)}{A(y, 2)} - E^2(y).
\end{aligned}$$

The lemma is proved. □

LEMMA 2.3.5.

$$D_x(A(y)) = 1 - \frac{A(y)}{A(y, 2)}.$$

PROOF.

$$\begin{aligned}
D_x(A(y)) &= \sum_{j=0}^r x_j \frac{\partial A(y)}{\partial x_j} \\
&= \sum_{j=0}^r x_j \sum_{i=0}^r \frac{\partial A(y)}{\partial y_i} \frac{\partial y_i}{\partial x_j} \\
&= \sum_{i=0}^r \frac{\partial A(y)}{\partial y_i} u_i \\
&= \sum_{i=0}^r -\frac{1}{(1+y_i)^2} \frac{y_i(1+y_i)}{1+2y_i} \frac{1}{A(y, 2)} \\
&= \frac{1}{A(y, 2)} \sum_{i=0}^r -\frac{y_i}{(1+y_i)(1+2y_i)} \\
&= \frac{1}{A(y, 2)} \sum_{i=0}^r \left(\frac{y_i}{1+y_i} - \frac{2y_i}{1+2y_i} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{A(y, 2)} (A(y, 2) - A(y)) \\
&= 1 - \frac{A(y)}{A(y, 2)}.
\end{aligned}$$

□

LEMMA 2.3.6.

$$(D_x + I_x)(g_1) = E^2(y)$$

PROOF.

$$\begin{aligned}
(D_x + I_x)(g_1) &= (D_x + I_x)(E^2(y)A(y)) \\
&= E^2(y)D_x(A(y)) + A(y)D_x(E^2(y)) + E^2(y)A(y) \\
&= E^2(y) \left(1 - \frac{A(y)}{A(y, 2)}\right) + A(y) \left(\frac{E^2(y)}{A(y, 2)} - E^2(y)\right) + E^2(y)A(y) \\
&= E^2(y).
\end{aligned}$$

□

Now from Lemma 2.3.4 to Lemma 2.3.6, we see that

$$\frac{E^2(y)}{A(y, 2)} = (D_x + I_x)^2(g_1).$$

LEMMA 2.3.7.

$$[x^k] \frac{E^2(y)}{A(y, 2)} = \prod_{i=0}^r \frac{n}{n - k_i} \binom{2n - 1 - k_i}{k_i}$$

where $n = k_0 + \cdots + k_r + 1$.

PROOF. Using multivariable Lagrange inversion (Lemma 1.2.1), we set

$$\phi_i = \frac{E^2(y)}{1 + y_i}.$$

Thus $y_i = x_i \phi_i(y)$. Using Lemma 1.2.2 to evaluate the determinant, we have

$$\begin{aligned}
[x^k] \frac{E^2(y)}{A(y, 2)} &= [y^k] \left\{ \frac{E^2(y)}{A(y, 2)} \phi^k(y) \left| \delta_{ij} - \frac{y_j}{\phi_i(y)} \frac{\partial \phi_i(y)}{\partial y_j} \right| \right\} \\
&= [y^k] \left\{ \frac{E^2(y)}{A(y, 2)} \phi^k(y) \frac{|\delta_{ij} + 2(\delta_{ij} - 1)y_j|}{E(y)} \right\} \\
&= [y^k] \left\{ \frac{E^2(y)}{A(y, 2)} \phi^k(y) \frac{E(y, 2)A(y, 2)}{E(y)} \right\} \\
&= [y^k] \left\{ \frac{E^{2n-1}(y)}{\prod_i (1 + y_i)^{k_i}} E(y, 2) \right\} \\
&= \prod_i \left[\binom{2n-1-k_i}{k_i} + 2 \binom{2n-1-k_i}{k_i-1} \right] \\
&= \prod_i \frac{n}{n-k_i} \binom{2n-1-k_i}{k_i}.
\end{aligned}$$

□

Therefore, Theorem 2.3.1 is proved.

REMARK 2.3.8. When $r = 1$, we have $k_0 + k_1 = n - 1$. Then the number is

$$\begin{aligned}
\frac{1}{n^2} \frac{n}{n-k_0} \binom{2n-1-k_0}{k_0} \frac{n}{n-k_1} \binom{2n-1-k_1}{k_1} \\
= \frac{(n+k_1)! (2n-1-k_1)!}{(k_1+1)! (2k_1+1)! (n-k_1)! (2n-1-2k_1)!},
\end{aligned}$$

which was found and proved by Bóna [3], Bousquet-Mélou [6], and Travis [25].

2.4. The number of two-stack-sortable r -permutations

The following theorem is a special case of Theorem 2.3.1 when we do not keep track of descents.

THEOREM 2.4.1. *The number of two-stack-sortable r -permutations of $[n]$ is:*

$$2(r+1) \frac{((2r+1)n)!}{n! (2rn+2)!}.$$

First proof:

We know that the generating function for two-stack-sortable r -permutations is given by (2.3.2). Setting $y_i = y$ for $i = 0, \dots, r$, which is equivalent to setting $x_i = x$ for $i = 0, \dots, r$, we get that

$$x = \frac{y}{(1+y)^{2r+1}}$$

and

$$\begin{aligned} g_1 &= (1+y)^{2r+2} \left(1 - (r+1) \frac{y}{1+y} \right) \\ &= (r+1)(1+y)^{2r+1} - r(1+y)^{2r+2} \\ &= (1+y)^{2r+1} (1 - ry) \\ (2.4.1) \quad &= \frac{y(1-ry)}{x}. \end{aligned}$$

Using Lagrange inversion [10, p. 17; 9], we set $\phi(y) = (1+y)^{2r+1}$. Then y satisfies the condition that $y = x\phi(y)$. Therefore,

$$\begin{aligned} [x^k]g_1 &= [x^{k+1}]xg_1 \\ &= \frac{1}{k+1} [y^k] \left\{ \frac{d(xg_1)}{dy} \phi^{k+1}(y) \right\} \\ &= \frac{1}{k+1} [y^k] \left\{ (1-2ry)(1+y)^{(2r+1)(k+1)} \right\} \\ &= \frac{1}{k+1} \left\{ \binom{(2r+1)(k+1)}{k} - 2r \binom{(2r+1)(k+1)}{k-1} \right\} \\ &= 2(r+1) \frac{((2r+1)(k+1))!}{(k+1)! (2r(k+1)+2)!}. \end{aligned}$$

Setting $n = k+1$, then the theorem is proved.

Second proof:

From Theorem 2.3.1, we know that the total number of two-stack-sortable permutations is

$$\frac{1}{n^2} \sum_{k_0 + \dots + k_r = n-1} \prod_{i=0}^r \frac{n}{n - k_i} \binom{2n-1-k_i}{k_i}.$$

Also we know that

$$(2.4.2) \quad c^n(t) = \sum_{i=0}^{\infty} \frac{n}{2i+n} \binom{2i+n}{i} t^i$$

for any integer n [19, p. 154]. Therefore,

$$\begin{aligned} [t^{k_i}] c^{-2n}(-t) &= [t^{k_i}] \sum_{j=0}^{\infty} \frac{2n}{2n-2j} \binom{2j-2n}{j} (-t)^j \\ &= \frac{2n}{2n-2k_i} (-1)^{k_i} \binom{2k_i-2n}{k_i} \\ &= \frac{n}{n-k_i} \binom{2n-1-k_i}{k_i}. \end{aligned}$$

Hence,

$$\begin{aligned} &\frac{1}{n^2} \sum_{k_0 + \dots + k_r = n-1} \prod_{i=0}^r \frac{n}{n - k_i} \binom{2n-1-k_i}{k_i} \\ &= \frac{1}{n^2} [t^{n-1}] (c^{2n}(-t))^{r+1} \\ &= \frac{1}{n^2} [t^{n-1}] \sum_{j=0}^{\infty} \frac{2n(r+1)}{2n(r+1)-2j} \binom{2j-2n(r+1)}{j} (-t)^j \\ &= \frac{1}{n^2} \frac{2n(r+1)}{2n(r+1)-2(n-1)} (-1)^{n-1} \binom{2(n-1)-2n(r+1)}{n-1} \\ &= \frac{r+1}{n(nr+1)} \binom{(2r+1)n}{n-1} \\ &= 2(r+1) \frac{((2r+1)n)!}{n! (2rn+2)!}. \end{aligned}$$

The theorem is proved. $\square$

REMARK 2.4.2. When $r = 1$, this number is

$$4 \frac{(3n)!}{n! (2n+2)!} = \frac{2(3n)!}{(n+1)! (2n+1)!},$$

which was conjectured by West [26, 27] and first proved by Zeilberger [28].

CHAPTER 3

Parallel results

3.1. Another approach

In the previous chapter, we counted the number of two-stack-sortable r -permutations with all types of descents. Here we approach a special case of the problem from another angle. We now count the number of two-stack-sortable r -permutations with only two basic parameters, the number of different letters and the number of components. We use the indeterminates x and $\bar{z}$ to count these two parameters respectively. The difference is, that *any component is allowed to be empty*.

Therefore, using the same decomposition method as in Chapter 2, if we let $p_n^{(k)}$ be the weight of all k -tuple r -permutations $(\alpha_1, \dots, \alpha_k)$ of $[n]$ where each of α_i ($i = 1, \dots, k$) is *allowed to be empty*, and let $q_n^{(k)} = p_n^{(r+k+1)}$,

$$(3.1.1) \quad p_k = \sum_{n=0}^{\infty} p_n^{(k)} x^n, \quad P = \sum_{k=0}^{\infty} p_k \bar{z}^k,$$

$$(3.1.2) \quad q_k = \sum_{n=0}^{\infty} q_n^{(k)} x^n, \quad Q = \sum_{k=0}^{\infty} q_k \bar{z}^k,$$

then analogous to Theorem 2.1.2, we have that the formal power series P and Q satisfy the functional equations

$$(3.1.3) \quad q_k = p_{r+k+1},$$

$$(3.1.4) \quad P = \frac{1}{1 - \bar{z}} + x\bar{z}PQ,$$

Since

$$Q = \frac{1}{\bar{z}^{r+1}} \left(P - \sum_{i=0}^r p_i \bar{z}^i \right),$$

we have that

THEOREM 3.1.1. *The generating function P of two-stack-sortable k -tuple r -permutations satisfies*

$$(3.1.5) \quad P = \frac{1}{1 - \bar{z}} + \frac{xP}{\bar{z}^r} \left(P - \sum_{i=0}^r p_i \bar{z}^i \right)$$

and P is uniquely determined as a power series by this functional equation.

Now, compared to the functional equations (2.1.10) and (2.1.11), this one is structurally simpler. We will also see that (3.1.5) can be easily derived from the functional equations (2.1.10) and (2.1.11).

If we set $x_i = x$ in functional equations (2.1.10) and (2.1.11), then they become

$$(3.1.6) \quad f_k = \sum_{j=0}^{r+1} \binom{r+1}{j} g_{k+j} x^j \text{ for } k \geq 0,$$

$$(3.1.7) \quad G = 1 + zFG.$$

Notice that p_k is the weight of of all k -tuple r -permutations where *any component is allowed to be empty* and g_k is the weight of of all k -tuple r -permutations where *no component is allowed to be empty*. Therefore,

$$(3.1.8) \quad p_k = \sum_{i=0}^k \binom{k}{i} g_i x^i,$$

which is equivalent to

$$P(x, \bar{z}) = \frac{1}{1 - \bar{z}} G \left(x, \frac{x\bar{z}}{1 - \bar{z}} \right).$$

Setting $x_i = x$, which is equivalent to setting $y_i = y$, in the solution of functional equations (Theorem 2.2.1) gives the solution to functional equation (3.1.6) and (3.1.7),

$$G = \frac{c(t)(1+y)^{r+1}}{(1+yc(t))^{r+1}},$$

where y is uniquely determined as formal power series in x by

$$x = \frac{y}{(1+y)^{2r+1}}$$

and $t = z(1+y)^{2(r+1)}$. Therefore,

THEOREM 3.1.2. *The solution to function equation (3.1.5) is:*

$$P = \left(1 + \frac{t}{y(1+y)}\right) \frac{c(t)(1+y)^{r+1}}{(1+yc(t))^{r+1}},$$

where y is uniquely determined as a formal power series in x by

$$x = \frac{y}{(1+y)^{2r+1}}$$

and $t = y(1+y)\frac{\bar{z}}{1-\bar{z}}$.

COROLLARY 3.1.3. *p_k is a polynomial in y with degree no greater than $2k$.*

PROOF.

$$\begin{aligned} x^k g_k &= x^k [z^k] G \\ &= \left[\left(\frac{z}{x} \right)^k \right] G \\ &= \left[\left(\frac{z}{x} \right)^k \right] \left\{ \frac{c(t)(1+y)^{r+1}}{(1+yc(t))^{r+1}} \right\} \\ &= [z^k] \left\{ \frac{c(xt)(1+y)^{r+1}}{(1+yc(xt))^{r+1}} \right\} \\ &= [z^k] \left\{ (1+y)^{r+1} \sum_{i=0}^{\infty} (-1)^i \binom{r+i}{i} y^i (c(xt))^{i+1} \right\} \end{aligned}$$

$$\begin{aligned}
&= [z^k] \left\{ (1+y)^{r+1} \sum_{i=0}^{\infty} (-1)^i \binom{r+i}{i} y^i \sum_{j=0}^{\infty} \frac{i+1}{2j+i+1} \binom{2j+i+1}{j} (xt)^j \right\} \\
&= [z^k] \left\{ (1+y)^{r+1} \sum_{i=0}^{\infty} (-1)^i \binom{r+i}{i} y^i \sum_{j=0}^{\infty} \frac{i+1}{2j+i+1} \binom{2j+i+1}{j} y^j (1+y)^j z^j \right\} \\
&= (1+y)^{r+k+1} y^k \sum_{i=0}^{\infty} (-1)^i \binom{r+i}{i} y^i \frac{i+1}{2k+i+1} \binom{2k+i+1}{k}.
\end{aligned}$$

The sum is a hypergeometric series that can be transformed by Euler's transformation [1]. Thus,

$$x^k g_k = y^k \sum_{i=0}^k (-1)^i \frac{(2i+1)r - (k-i)}{(r-k)(k+i+1)} \binom{r-k}{i} \binom{2k}{k-i} y^i.$$

Therefore, $x^k g_k$ is a polynomial in y with degree no greater than $2k$. Thus p_k is a polynomial in y with degree no greater than $2k$ because $p_k = \sum_{i=0}^k \binom{k}{i} x^i g_i$ (Eq. (3.1.8)).

□

In particular,

COROLLARY 3.1.4. p_1 is the generating function of two-stack-sortable r -permutations, and

$$p_1 = 1 + y - ry^2.$$

PROOF. Since $p_1 = 1 + xg_1$ (Eq. (3.1.8)) and $g_1 = \frac{y(1-ry)}{x}$ (Eq. (2.4.1)),

$$\begin{aligned}
p_1 &= 1 + xg_1 \\
&= 1 + y - ry^2.
\end{aligned}$$

□

3.2. Connection to Zeilberger's functional equation

Zeilberger first proved West's conjecture that the number of two-stack-sortable permutations of length n is $\frac{2(3n)!}{(n+1)!(2n+1)!}$ [28]. He used a factorization similar to ours and the functional equation he got is equivalent to our functional equation (3.1.5) in the case of $r = 1$.

In Zeilberger's paper [28], he defined $i(\pi)$ (where π is any permutation of $\{1, 2, \dots, n\}$) to be the largest integer i such that the subsequence of the 'big i ': $\{n-i+1, \dots, n-1, n\}$ are in decreasing order, defined $W^{(i)}$ to be the set of all permutations (of any length) π such that $i(\pi) = i$, and let $W^{\geq i}$ to be the set of all permutations π such that $i(\pi) \geq i$. Also he defined $W^{(i)}(x)$ to be the formal power series that equals to the sum of all the weights of elements of $W^{(i)}$, and $W^{\geq i}(x)$ to be the formal power series that equals to the sum of all the weights of elements of $W^{\geq i}$, and he defined

$$\begin{aligned}\Phi(x, t) &:= \sum_{i=0}^{\infty} W^{(i)}(x) t^i, \\ \bar{\Phi}(x, t) &:= \sum_{i=0}^{\infty} W^{\geq i}(x) t^i.\end{aligned}$$

Then, he got

$$(3.2.1) \quad \Phi(x, t) = \frac{1}{1 - xt} + \frac{xt(\Phi(x, 1) - t\Phi(x, t))(\Phi(x, 1) - \Phi(x, t))}{(1 - t)^2}.$$

Now noticing that $\Phi(x, 1) = \bar{\Phi}(x, 0)$ and

$$\bar{\Phi}(x, t) = \frac{\Phi(x, 1) - t\Phi(x, t)}{1 - t}.$$

Equation (3.2.1) is equivalent to

$$\bar{\Phi}(x, t) = \frac{1}{1 - xt} + \frac{(1 + xt\bar{\Phi}(x, t))(\bar{\Phi}(x, t) - \bar{\Phi}(x, 0))}{t}.$$

Now it is easy to check that $P(x, \bar{z}) = 1 + \bar{z}\bar{\Phi}\left(x, \frac{\bar{z}}{x}\right)$. Therefore, the functional equation (3.2.1) is equivalent to (3.1.5) in the case of $r = 1$.

The combinatorial connection between the set $W^{\geq i}$ and the set of k -tuple permutations was clearly stated in Zeilberger's paper [28]. For a typical member π of $W^{\geq i}$, if its length is n , then it has the form

$$\pi = \alpha_0 n \alpha_1 (n-1) \cdots (n-i+1) \alpha_i,$$

where $\alpha_0, \dots, \alpha_i$ are (possibly empty) permutations of disjoint smaller sets, the union of whose underlying sets is $\{1, 2, \dots, n-i\}$. Now, by iterating the definition of the stack sorting operation S ,

$$S(\pi) = S(\alpha_0)S(\alpha_1) \cdots S(\alpha_i)(n-i+1)(n-i+2) \cdots n,$$

so that,

$$S^2(\pi) = S(S(\alpha_0)S(\alpha_1) \cdots S(\alpha_i))(n-i+1)(n-i+2) \cdots n.$$

It follows that there is a 1-1 correspondence between the elements of $W^{\geq i}$ and $(i+1)$ -tuple permutations $\alpha_0, \alpha_1, \dots, \alpha_i$, such that $S(S(\alpha_0) \cdots S(\alpha_i)) = I$, and the underlying sets of the α 's are disjoint and their union is $\{1, 2, \dots, n-i\}$.

CHAPTER 4

Further results

4.1. A generalization of the stack-sorting operation on r -permutations

Here we introduce a more general form of the stack-sorting operation on r -permutations.

DEFINITION 4.1.1. For a given positive number r , let $\lambda = (\lambda_0, \lambda_1, \dots, \lambda_l)$ satisfy the condition that $0 < \lambda_0 < \lambda_1 < \dots < \lambda_l = r$. Let π be an r -permutation of $[n]$. Then we can write π as $\pi = \alpha_0 n \alpha_1 n \dots n \alpha_r$. The generalized stack-sorting operation S_λ on r -permutations is defined by

$$S_\lambda(\pi) = S_\lambda(\alpha_0) \cdots S_\lambda(\alpha_{\lambda_0}) n S_\lambda(\alpha_{\lambda_0+1}) \cdots S_\lambda(\alpha_{\lambda_1}) n \cdots n S_\lambda(\alpha_{\lambda_{l-1}+1}) \cdots S_\lambda(\alpha_{\lambda_l}) n.$$

Notice that $S_\lambda(\pi)$ is an $(l+1)$ -permutation and $S_\lambda = S$ when $l = 0$, where S is the ordinary stack sorting operation on r -permutations (Definition 1.1.2).

It is clear that it is not interesting to consider stack-sortable r -permutation under this definition. Also, in order to consider two-stack-sortable r -permutations under this definition, we have to consider $S(S_\lambda(\pi))$ instead of $S_\lambda^2(\pi)$.

To enumerate the number of two-stack-sortable r -permutations with descents under this definition of stack-sorting operation, we still weight any type i descent by x_i . Furthermore, we denote $x^{(0)} = (x_0, x_1, \dots, x_{\lambda_0})$ and $x^{(i)} = (x_{\lambda_{i-1}+1}, \dots, x_{\lambda_i})$ for $1 \leq i \leq l$. Also, we use u to keep track of the number of different letters.

Now let $g_n^{(k)}$ be the sum of the weights of two-stack-sortable k -tuple r -permutations $(\alpha_1, \alpha_2, \dots, \alpha_k)$ such that every α_i is nonempty and let

$$f_{n,0}^{(k)} = \sum_{j=0}^{\lambda_0+1} g_n^{(k+j)} e_j(x^{(0)}),$$

$$f_{n,i}^{(k)} = \sum_{j=0}^{\lambda_i - \lambda_{i-1}} g_n^{(k+j)} e_j(x^{(i)}) \text{ for } 1 \leq i \leq l.$$

Also let

$$(4.1.1) \quad \begin{aligned} f_{k,i} &= \sum_{n=0}^{\infty} f_{n,i}^{(k)} u^n, & F_i &= \sum_{k=0}^{\infty} f_{k,i} z^k, \\ g_k &= \sum_{n=0}^{\infty} g_n^{(k)} u^n, & G &= \sum_{k=0}^{\infty} g_k z^k. \end{aligned}$$

Then we have the following functional equations by the same reasoning as in section 2.1:

LEMMA 4.1.2. *Let G and $f_{k,i}$ be defined as above, then*

$$(4.1.2) \quad G = 1 + zuGF_0 \prod_{i=1}^l f_{0,i},$$

$$(4.1.3) \quad f_{k,0} = \sum_{j=0}^{\lambda_0+1} g_{k+j} e_j(x^{(0)}),$$

$$(4.1.4) \quad f_{k,i} = \sum_{j=0}^{\lambda_i - \lambda_{i-1}} g_{k+j} e_j(x^{(i)}) \text{ for } 1 \leq i \leq l.$$

If we let $A = \prod_{i=1}^l f_{0,i}$, we can make the substitution $x = uA$ so that the functional equations (4.1.2) to (4.1.4) become

$$(4.1.5) \quad G = 1 + zxGF_0,$$

$$(4.1.6) \quad f_{k,0} = \sum_{j=0}^{\lambda_0+1} g_{k+j} e_j(x^{(0)}).$$

By Theorem 2.2.1, the solution to these functional equations is:

$$G = \frac{c(t)E(y)}{E(y, c(t))},$$

$$F_0 = \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right),$$

where $y = (y_0, \dots, y_{\lambda_0})$ is uniquely determined by $xx_j = \frac{y_j(1+y_j)}{E^2(y)}$ for $0 \leq j \leq \lambda_0$, and $t = zxE^2(y)$. From the solution, we can express g_i in terms of x and y , i.e., $g_i = g_i(x, y)$. Since $x = uA$, g_i can be expressed in terms of u, A, y , i.e., $g_i = g_i(u, A, y)$. By equation (4.1.4),

$$A = \prod_{i=1}^l f_{0,i}$$

$$= \prod_{i=1}^l \sum_{j=0}^{\lambda_i - \lambda_{i-1}} g_j(u, A, y) e_j(x^{(i)}).$$

Now we can solve for A from the equation above so that A is expressed in terms of $u, y, x^{(i)}$, $1 \leq i \leq l$, i.e., $A = A(u; y; x^{(1)}; \dots; x^{(l)})$. Therefore, given $g_i = g_i(u, A, y)$ and $x_j = \frac{y_j(1+y_j)}{xE^2(y)}$ for $0 \leq j \leq \lambda_0$, we can solve for g_i (for example, using Lagrange inversion) so that $g_i = g_i(u; x^{(0)}; x^{(1)}; \dots; x^{(l)})$.

Therefore,

THEOREM 4.1.3. *If G and F_i are defined in (4.1.1), then G and F_i satisfy the following functional equations:*

$$G = \frac{c(t)E(y)}{E(y, c(t))},$$

$$F_0 = \frac{E(y)}{t} \left(E(y) - \frac{E(y, c(t))}{c(t)} \right),$$

where $y = (y_0, \dots, y_{\lambda_0})$ is uniquely determined by $xx_i = \frac{y_i(1+y_i)}{E^2(y)}$ for $0 \leq i \leq \lambda_0$, and $t = zxE^2(y)$. In particular, the g_i are algebraic.

REMARK 4.1.4. It is easy to get $g_0 = 1$. Also, similar to Equation (2.3.2), we can get

$$g_1 = xE(y) \sum_{n=0}^{\infty} (1-n)e_n(y)$$

in this case.

For example, the simplest nontrivial case is when $l = 1$ and $\lambda = (1, 2)$. We have $x^{(0)} = (x_0, x_1)$, $x^{(1)} = (x_2)$, and the functional equations for this case are:

$$G = 1 + zuGF_0f_{0,1}$$

$$f_{k,0} = g_k + g_{k+1}(x_0 + x_1) + g_{k+2}x_0x_1$$

$$f_{k,1} = g_k + g_{k+1}x_2$$

Now we make the substitution $x = uf_{0,1}$. Then the functional equations above become

$$G = 1 + zxGF_0$$

$$f_{k,0} = g_k + g_{k+1}(x_0 + x_1) + g_{k+2}x_0x_1$$

By Theorem 3.1.2, the solution to these functional equations is:

$$G = \frac{c(t)E(y)}{E(y, c(t))},$$

where $y = (y_0, y_1)$, $E(y) = (1 + y_0)(1 + y_1)$ and y is uniquely determined by $xx_i = \frac{y_i(1 + y_i)}{E^2(y)}$ for $i = 0, 1$, and $t = zx E^2(y)$.

Also, from Remark 4.1.4, we have

$$\begin{aligned}
 (4.1.7) \quad & g_0 = 1, \\
 & g_1 = xE(y) \sum_{n=0}^{\infty} (1-n)e_n(y) \\
 & = x(1+y_0)(1+y_1)(1-y_0y_1) \\
 & = uf_{0,1}(1+y_0)(1+y_1)(1-y_0y_1).
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 f_{0,1} &= g_0 + g_1x_2 \\
 &= 1 + uf_{0,1}(1+y_0)(1+y_1)(1-y_0y_1)x_2.
 \end{aligned}$$

By solving for $f_{0,1}$, we get

$$f_{0,1} = \frac{1}{1 - u(1+y_0)(1+y_1)(1-y_0y_1)x_2}.$$

So, by (4.1.7),

$$g_1 = \frac{u(1+y_0)(1+y_1)(1-y_0y_1)}{1 - u(1+y_0)(1+y_1)(1-y_0y_1)x_2}.$$

Now, since $xx_i = \frac{y_i(1+y_i)}{E^2(y)}$ for $i = 0, 1$ and $x = uf_{0,1}$, we have

$$ux_i = \frac{y_i(1+y_i)(1 - u(1+y_0)(1+y_1)(1-y_0y_1)x_2)}{E^2(y)} \text{ for } i = 0, 1.$$

Since (y_0, y_1) is uniquely determined by the equation above, we can solve for g_1 in terms of u, x_0, x_1 and x_2 by using Lagrange inversion.

We can also approach the problem without counting descents in the same way as what we did in Section 3.1. If we let P be the generating function of two-stack-sortable k -tuple r -permutations under S_λ for any k , with u and $\bar{z}$ keeping track of the number of different letters and the number of components, then

LEMMA 4.1.5. *P satisfies*

$$(4.1.8) \quad P = \frac{1}{1 - \bar{z}} + \frac{uP}{\bar{z}^{\lambda_0}} \left(P - \sum_{i=0}^{\lambda_0} p_i \bar{z}^i \right) \prod_{i=1}^l p_{\lambda_i - \lambda_{i-1}}.$$

To solve this functional equation, first we make the substitution

$$x = u \prod_{i=1}^l p_{\lambda_i - \lambda_{i-1}}.$$

Then (4.1.8) becomes

$$P = \frac{1}{1 - \bar{z}} + \frac{xP}{\bar{z}^{\lambda_0}} \left(P - \sum_{i=0}^{\lambda_0} p_i \bar{z}^i \right).$$

From Theorem 3.1.2, we know that

THEOREM 4.1.6. *The solution to the functional equation above is*

$$P = \left(1 + \frac{t}{y(1+y)} \right) \frac{c(t)(1+y)^{\lambda_0+1}}{(1+yc(t))^{\lambda_0+1}},$$

where y is uniquely determined as a formal power series in x by

$$x = \frac{y}{(1+y)^{2\lambda_0+1}},$$

$$\begin{aligned} t &= x(1+y)^{2(\lambda_0+1)} \frac{\bar{z}}{1-\bar{z}} \\ &= y(1+y) \frac{\bar{z}}{1-\bar{z}} \end{aligned}$$

and

$$x = u \prod_{i=1}^l p_{\lambda_i - \lambda_{i-1}}.$$

Since p_k is a polynomial in y (Theorem 3.1.3) and

$$u = \frac{y}{(1+y)^{2\lambda_0+1} \prod_{i=1}^l p_{\lambda_i - \lambda_{i-1}}},$$

we can solve for p_k in terms of u .

Again, the simplest nontrivial case is when $l = 1$ and $\lambda = (1, 2)$, and the functional equation for this case is:

$$P = \frac{1}{1 - \bar{z}} + up_1 \frac{P(P - p_0 - p_1 \bar{z})}{\bar{z}}.$$

Now make the substitution $x = up_1$. Then the functional equation above becomes

$$P = \frac{1}{1 - \bar{z}} + x \frac{P(P - p_0 - p_1 \bar{z})}{\bar{z}}.$$

From Corollary 3.1.4, $p_1 = 1 + y - y^2$, where y is uniquely determined by $x = \frac{y}{(1+y)^3}$,

or

$$u = \frac{y}{(1+y)^3(1+y-y^2)}.$$

Now we can use Lagrange inversion [10, p. 17; 9] to solve for p_1 .

Since $y = u(1+y)^3(1+y-y^2)$,

$$\begin{aligned} [u^n]p_1 &= \frac{1}{n}[y^{n-1}] \frac{dp_1}{dy} (1+y)^{3n} (1+y-y^2)^n \\ &= \frac{1}{n}[y^{n-1}] (1-2y)(1+y)^{3n} \sum_{i=0}^n (-1)^i \binom{n}{i} (1+y)^i y^{2(n-i)} \\ &= \frac{1}{n}[y^{n-1}] (1-2y) \sum_{i=0}^n \sum_{j=0}^{3n+i} (-1)^i \binom{n}{i} \binom{3n+i}{j} y^j y^{2(n-i)} \\ &= \sum_{i=0}^n (-1)^{n-i} \frac{1}{n} \binom{n}{i} \left[\binom{3n+i}{2i-1-n} - 2 \binom{3n+i}{2i-2-n} \right]. \end{aligned}$$

4.2. A characterization of t -stack-sortable permutations

It is natural to consider counting k -stack-sortable permutations for $k > 2$ now. But although people have been trying, little has been found yet, even for three-stack-sortable permutations.

Similar to West's characterization of two-stack-sortable permutations (which did not lead to an enumeration) [26, 27], we can give a characterization for t -stack-sortable permutations.

For a sequence $a_1 a_2 \cdots a_n$ of different letters, define (a_i, a_j) to be an inversion of the sequence if $1 < i < j < n$ and $a_i > a_j$. We also define $\text{rank}(a_i)$ as follows: $\text{rank}(a_i) = m$ if there are exactly $m - 1$ letters among $a_1, a_2, \dots, a_n$ that are smaller than a_i .

THEOREM 4.2.1. *A permutation π is t -stack-sortable if and only if it does not contain a subsequence $\pi' = a_1 a_2 \cdots a_{t+2}$ which satisfies the following conditions:*

- (1) $\text{rank}(a_{t+2}) = 1$;
- (2) $\text{rank}(a_{t+1}) = t + 2$;
- (3) *For any i and j such that $1 \leq i < j \leq t$ and (a_i, a_j) is an inversion, there does not exist a subsequence $\pi'' = c_1 c_2 \cdots c_s$ of π , where $a_i < c_1 < c_2 < \cdots < c_s$ and $s = t + 2 - \text{rank}(a_i)$, such that π'' appears between a_i and a_j in π .*

To prove this theorem, we need some lemmas from West [26, 27].

LEMMA 4.2.2. *(West) If π is a permutation of $[n]$ and $1 \leq a < b \leq n$, and if a precedes b in π , then a precedes b in $S(\pi)$.*

LEMMA 4.2.3. (West) *If π is a permutation of $[n]$ and $1 \leq a < b \leq n$, and if b precedes a in π , then b precedes a in $S(\pi)$ if there exists $c > b$ such that b precedes c and c precedes a in π . If there is no such c , then a precedes b in $S(\pi)$.*

LEMMA 4.2.4. (West) *If b and a form an inversion in $S(\pi)$, then there exist $c > b$ such that b precedes c and c precedes a in π .*

Now let's prove Theorem 4.2.1.

PROOF. Tarjan and West proved that the theorem is correct when $t = 1$ and $t = 2$ [24, 26, 27]. We now prove the theorem by induction.

Suppose a permutation π contains a subsequence $\pi' = a_1 a_2 \cdots a_{t+2}$ satisfying the conditions (1) – (3). Let $\pi_1 = S(\pi)$ and suppose that $b_1 b_2 \cdots b_{t+2}$ is the subsequence of π_1 that was π' in π .

Since π' satisfies conditions (1) – (2), by Lemma 4.2.2, a_i precedes a_{t+1} in $S(\pi)$ for $1 \leq i \leq t$; by Lemma 4.2.3, a_i precedes a_{t+2} in $S(\pi)$ for $1 \leq i \leq t$. Thus, we get either $\text{rank}(b_{t+2}) = t + 2$, $\text{rank}(b_{t+1}) = 1$ or $\text{rank}(b_{t+2}) = 1$, $\text{rank}(b_{t+1}) = t + 2$. Let $\pi'_1 = b_1 b_2 \cdots b_t b_{t+1}$ by supposing $\text{rank}(b_{t+1}) = 1$ (If $\text{rank}(b_{t+2}) = 1$, we let $\pi'_1 = b_1 b_2 \cdots b_t b_{t+2}$). Therefore, π'_1 satisfies condition (1) with t replaced by $t - 1$.

Let $\text{rank}(a_{i_0}) = t + 1$. Then $a_{i_0} > a_i$ for $1 \leq i \leq t$ and $i \neq i_0$. By Lemma 4.2.2, a_i precedes a_{i_0} in π_1 for $1 \leq i < i_0$. If $i_0 \neq t$, then a_{i_0} and a_i form an inversion in π' for $i_0 < i \leq t$. Since $t + 2 - \text{rank}(a_{i_0}) = 1$, by condition (3), there is no letter bigger than a_{i_0} appears between a_{i_0} and a_i in π . By Lemma 4.2.3, a_i precedes a_{i_0} in π_1 for $i_0 < i \leq t$. Therefore, a_i precedes a_{i_0} in π_1 for $1 \leq i \leq t$ and $i \neq i_0$. So $b_t = a_{i_0}$ and $\text{rank}(b_t) = t + 1$. Hence π'_1 satisfies condition (2).

Suppose that π'_1 does not satisfy condition (3); that is, there exist $1 \leq i < j \leq t - 1$ such that $b_i > b_j$, and there exists a subsequence $\pi''_1 = c_1 c_2 \cdots c_s$ where $b_i < c_1 < c_2 <$

$\cdots < c_s$ and $s = t + 1 - \text{rank}(b_i)$, such that π_1'' appears between b_i and b_j in π_1 . Then c_s and b_j form an inversion in π_1 . By Lemma 4.2.4, there exists some $c > c_s$ such that c_s precedes c and c precedes b_j in π . This contradicts the fact that π' satisfies condition (3).

Now, since π_1' satisfies condition (1) – (3), by the induction hypothesis, π_1 is not $(t - 1)$ -stack-sortable. So π is not t -stack-sortable.

Conversely, we can show that if π fails to be t -stack-sortable, then it contains a subsequence that satisfies the three conditions.

If π is not t -stack-sortable, then $\pi_1 = S(\pi)$ is not $(t - 1)$ -stack-sortable. By the induction hypothesis, π_1 contains a subsequence $\pi_1' = b_1 b_2 \cdots b_{t+1}$ which satisfies the following conditions:

- (1) $\text{rank}(b_{t+1}) = 1$;
- (2) $\text{rank}(b_t) = t + 1$;
- (3) For any i and j such that $1 \leq i < j \leq t - 1$ and $b_i > b_j$, there does not exist a subsequence $\pi_1'' = c_1 c_2 \cdots c_s$ where $b_i < c_1 < c_2 < \cdots < c_s$ and $s = t + 1 - \text{rank}(b_i)$, such that π_1'' appears between b_i and b_j in π_1 .

Notice that b_i and b_{t+1} form an inversion in π_1 for $1 \leq i \leq t$. By Lemma 4.2.4, b_i precedes b_{t+1} in π for $1 \leq i \leq t$. In particular, there exists some $a > b_t$ such that b_i precedes a and a precedes b_{t+1} in π for $1 \leq i \leq t$. Suppose that $a_1 a_2 \cdots a_{t+1}$ is the subsequence of π that gets transformed to π_1' . Then it is clear that $a_{t+1} = b_{t+1}$. Let $\pi' = a_1 a_2 \cdots a_t a a_{t+1}$. Then π' satisfies conditions (1) and (2).

If π' does not satisfy condition (3); that is, there exist $1 \leq i < j \leq t$ such that $a_i > a_j$, and there exists a subsequence $\pi'' = c_1 c_2 \cdots c_s$ where $a_i < c_1 < c_2 < \cdots < c_s$ and $s = t + 2 - \text{rank}(a_i)$, such that π'' appears between a_i and a_j , then by Lemma

4.2.2, the subsequence $c_1c_2\ldots c_{s-1}$ appears between a_i and a_j in π_1 . This contradicts the fact that π'_1 satisfies condition (3). Therefore π' satisfies condition (3).

The theorem is proved. $\square$

LEMMA 4.2.5. (*Tarjan*) *A permutation π is stack-sortable if and only if π contains no subsequence of type 231.*

PROOF. This lemma is the case $t = 1$ of Theorem 4.2.1.

By Theorem 4.2.1, a permutation π is stack-sortable if and only if it does not contain a subsequence $\pi' = a_1a_2a_3$, which satisfies the conditions that $\text{rank}(a_3) = 1$ and $\text{rank}(a_2) = 3$, which means that $a_3 < a_1 < a_2$. Therefore, π' is a type 231 subsequence. $\square$

LEMMA 4.2.6. (*West*) *A permutation fails to be two-stack-sortable if it contains a subsequence of type 2341 or a subsequence of type 3241 which is not part of a subsequence of type 35241. If it contains no such subsequence, π is two-stack-sortable.*

PROOF. This lemma is the case $t = 2$ of Theorem 4.2.1.

By Theorem 4.2.1, a permutation π is two-stack-sortable if and only if it does not contain a subsequence $\pi' = a_1a_2a_3a_4$, which satisfies the conditions that (1) $\text{rank}(a_4) = 1$, (2) $\text{rank}(a_3) = 4$ and, (3) $a_1 < a_2$ or $a_1 > a_2$. If $a_1 < a_2$, then π' is a subsequence of type 2341. If $a_1 > a_2$, then π' is a subsequence of type 3241. By condition (3), there does not exist a $c > a_1$ appearing between a_1 and a_2 . If there does exist such a c , then when $c > a_3$, $a_1ca_2a_3a_4$ is a subsequence of type 35241; when $c < a_3$, $ca_2a_3a_4$ is a subsequence of 3241, which needs to be considered again in the same way. Therefore, these three conditions suggest that π' is a either type 2341 or type 3241 subsequence. If it is a type 3241 subsequence, then it is not part of a subsequence of type 35241. $\square$

To enumerate the three-stack-sortable permutations by the same decomposition we used for two-stack-sortable permutations, we consider the following object, called a μ -tuple permutation, where $\mu = (\mu_1, \dots, \mu_k)$ and μ_i are positive integers.

DEFINITION 4.2.7. $((\alpha_1, \dots, \alpha_{\mu_1}), (\alpha_{\mu_1+1}, \dots, \alpha_{\mu_1+\mu_2}), \dots, (\alpha_{\mu_1+\dots+\mu_{k-1}+1}, \dots, \alpha_{\mu_1+\dots+\mu_k}))$ is a μ -tuple permutation of $[n]$ if $\alpha_1 \cdots \alpha_{\mu_1+\dots+\mu_k}$ is a permutation of $[n]$. A μ -tuple permutation is three-stack-sortable if

$$S(S(S(\alpha_1) \cdots S(\alpha_{\mu_1}))S(S(\alpha_{\mu_1+1}) \cdots S(\alpha_{\mu_1+\mu_2})) \cdots S(S(\alpha_{\mu_1+\dots+\mu_{k-1}+1}) \cdots S(\alpha_{\mu_1+\dots+\mu_k}))) = I.$$

By the same reasoning as in the case of two-stack-sortable permutations, if g_μ is the generating function for three-stack-sortable μ -tuple permutations, we get

$$g_\mu = 1 + x \left(\sum_{i=0}^{\mu_k-1} g_{(\mu_1, \dots, \mu_k-i+1, i)} + \sum_{i=0}^{\mu_{k-1}-1} g_{(\mu_1, \dots, \mu_{k-1}-i+1, i)} g_{(\mu_k)} + \cdots + \sum_{i=0}^{\mu_1-1} g_{(\mu_1-i+1)} g_{(\mu_2, \dots, \mu_k)} \right).$$

It seems that this functional equation is hard to solve.

4.3. A modification on the functional equations.

If we replace the elementary symmetric functions $e_i(x)$ by complete homogeneous symmetric functions $h_i(x)$, then the functional equations in Theorem 2.1.2 will become:

$$f_k = \sum_{j=0}^{\infty} g_{k+j} h_j(x)$$

$$G = 1 + zFG$$

Combinatorially, in terms of the tree representation, instead of having at most one child of each type, now we can have any number of children of each type. To solve

this, everything else will be the same except one modification on the substitution

$$x_i = \frac{y_i(1 + y_i)}{E^2(y)},$$

which in this case will be

$$x_i = \frac{y_i(1 - y_i)}{H^2(y)},$$

where $H(y)$ is defined analogously to $E(y)$. Then analogous to the identity in Lemma 2.2.2, we get

$$H\left(x, \frac{1}{z}\right) = H(y, c(t))H\left(y, \frac{1}{tc(t)}\right)$$

The solutions to these functional equations will be:

$$G = \frac{c(t)H(y)}{H(y, c(t))},$$

$$F = \frac{H(y)}{t} \left(H(y) - \frac{H(y, c(t))}{c(t)} \right).$$

Therefore,

$$g_1 = f_0$$

$$= H(y) \sum_{n=0}^{\infty} (1 - n) h_n(y)$$

Again, we use multivariable Lagrange inversion to get the coefficient of x^k in g_1 , where $x = (x_0, \dots, x_r)$ and $k = (k_0, \dots, k_r)$ and it is

$$\frac{1}{n^2} \prod_{k=0}^r \frac{n}{n + k_i} \binom{2n + 2k_i}{k_i},$$

where $n = 1 + k_0 + k_1 + \dots + k_r$.

REMARK 4.3.1. If we let

$$A(n; k_0, \dots, k_r) = \frac{1}{n^2} \prod_{k=0}^r \frac{n}{n - k_i} \binom{2n - 1 - k_i}{k_i}$$

and

$$B(n; k_0, \dots, k_r) = \frac{1}{n^2} \prod_{k=0}^r \frac{n}{n+k_i} \binom{2n+2k_i}{k_i},$$

then

$$A(-n; k_0, \dots, k_r) = (-1)^{n-1} B(n; k_0, \dots, k_r).$$

Bibliography

- [1] W. N. Bailey, *Generalized Hypergeometric Series*, Cambridge, The University Press, 1935.
- [2] M. Bóna, Exact enumeration of 1342-avoiding permutations: a close link with labeled trees and planar maps, *J. Combin. Theory Ser. A* 80 (1997), 257–272.
- [3] M. Bóna, 2-stack sortable permutations with a given number of runs, MSRI Preprint 1997-055.
- [4] M. Bousquet-Mélou, Sorted and/or sortable permutations, *Discrete Math*, 225 (2000), 25–50.
- [5] M. Bousquet-Mélou, On (some) functional equations arising in enumerative combinatorics, preprint.
- [6] M. Bousquet-Mélou, Multi-statistic enumeration of two-stack sortable permutations, *Electron. J. Combin*, 5 (1998), no. 1, Research Paper 21, 12 pp.
- [7] S. Dulucq, S. Gire and O. Guibert, A combinatorial proof of J. West’s conjecture. *Discrete Math*, 187 (1998), 71–96.
- [8] Paul H. Edelman, Chain enumeration and noncrossing partitions, *Discrete Math*. 31 (1980), 171–180.
- [9] Ira M. Gessel, A combinatorial proof of the multivariable Lagrange inversion formula, *J. Combin. Theory Ser. A* 45 (1987), 178–195.
- [10] I. P. Goulden and D. M. Jackson, *Combinatorial Enumeration*, John Wiley and Sons, 1983.
- [11] I. P. Goulden and J. West, Raney paths and a combinatorial relationship between rooted nonseparable planar maps and two-stack-sortable permutations, *J. Combin. Theory Ser. A* 75 (1996), 220–242.
- [12] F. K. Hwang and C. L. Mallows, Enumerating nested and consecutive partitions, *J. Combin. Theory Ser. A* 70 (1995), 323–333.

- [13] Paul Klingsberg and Cynthia Schmalzried, Barred permutations, Proceedings of the Twenty-fourth Southeastern International Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1993), *Congr. Numer.*, 95 (1993), 153–161.
- [14] Paul Klingsberg and Cynthia Schmalzried, A family of constructive bijections involving Stirling permutations, Proceedings of the Twenty-first Southeastern Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1990), *Congr. Numer.*, 78 (1990), 11–15.
- [15] D.E. Knuth, *The Art of Computer Programming*, Vol. 1 Addison-Wesley, Reading, MA, 2nd ed., 1973.
- [16] SeungKyung Park, The r -multipermutations, *J. Combin. Theory Ser. A* 67 (1994), 44–71.
- [17] SeungKyung Park, P -partitions and q -Stirling numbers, *J. Combin. Theory Ser. A* 68 (1994), 33–52.
- [18] SeungKyung Park, Inverse descents of r -multipermutations, *Discrete Math.*, 132 (1994), 215–229.
- [19] John Riordan, *Combinatorial Identities*, Wiley, New York, 1968.
- [20] Rodica Simion, Noncrossing partitions, *Discrete Math.* 217 (2000), 367–409.
- [21] Richard P. Stanley, Parking functions and noncrossing partitions, The Wilf Festschrift (Philadelphia, PA, 1996), *Electron. J. Combin.* 4 (1997), no. 2.
- [22] Robert A. Sulanke, Refinements of the Narayana numbers, *Bull. Inst. Combin. Appl.* 7 (1993), 60–66.
- [23] Robberto Tomassia and Michael T. Goodrich, *Data Structures and Algorithms in Java*, second edition, John Wiley and Sons, 2001.
- [24] Robert Tarjan, Sorting using networks of queues and stacks, *J. Assoc. Comput. Mach.* 19 (1972), 341–346.
- [25] Leopold Travis, A proof of two conjectured refinements of West’s formula for the number of two-stack-sortable permutations, unpublished.
- [26] J. West, Permutations with forbidden subsequences; and, stack sortable permutations, Ph.D. thesis, MIT, 1990
- [27] J. West, Sorting twice through a stack, *Theoretical Computer Science*, 117 (1993) 303–313.
- [28] D. Zeilberger, A proof of Julian West’s conjecture that the number of two-stack-sortable permutations of length n is $2(3n)!/((n+1)!(2n+1)!)$, *Discrete Math.* 102 (1992), 85–93.